

Prevalence and patterns of smoking and smokeless tobacco use in Santa-Cruz, Goa, India.

Gaude Shital Rama¹, Silva Pinto Vanita²

¹Junior Consultant, NCDC, Directorate of Health Services, Campal, Panaji, Goa.

²Assistant Professor, Dept. Of Preventive and Social Medicine, Goa Medical College, Bambolim Goa.

(Received:05/12/2014, Revised: 01/02/2015, Accepted: 12/03/2015)

Abstract :

Background : Tobacco use is the single most preventable cause of premature disease and death, currently leading to over five million deaths globally each year (one of every six adult deaths) which is expected to rise to over eight million deaths by 2030.

Aims : This paper examines the prevalence of tobacco use in the individuals aged 20 years and above in the urban field practice area of Santa Cruz, the prevalence of the various forms of tobacco used and some socio-demographic factors associated with tobacco use.

Materials and Methods : The present cross sectional community based study was conducted in Santa Cruz an urban area of Goa. 311 households were chosen using systematic random sampling and 935 individuals aged 20 years and above from the 311 households were recruited in the study. The data was collected in a pre-designed structured questionnaire by face to face interview.

Results : The prevalence of tobacco use was found to be 34.8%. The prevalence of smoking tobacco use was found to be 20.3%, smokeless tobacco use was found to be 14.1 % and tobacco use in any form was found to be 0.4%. Use of cigarette (92%) was the most common form of smoking tobacco used and Betel Quid (51%) was the most common form of smokeless tobacco used. Tobacco use was found to be highest (40.8%) in the age group of 40-49 years. Tobacco use was higher in males as compared to females. Tobacco use was higher in illiterate (40%) , unemployed (65.2%) and low socio-economic status (51.5%). Mean age at initiation of tobacco use in males was 18.85 ± 4.3 years and in females the mean age at initiation was 18.19 ± 1.5 years.

Conclusion : Tobacco Control Policy would have to consider the type of tobacco consumption, age at initiation of tobacco use, education and employment status of individuals while striving to decrease tobacco use. Priority should also be given to smokeless tobacco control efforts.

Key Words : Smokeless Tobacco, Smoking, Tobacco use, Urban Population.

Introduction: Tobacco use is the single most preventable cause of premature disease, disability and death, currently leading to over five million deaths each year worldwide which is expected to rise to over eight million deaths, that is, one of every six adult deaths yearly by 2030. Giving the existing trends in mortality, about 500 million people alive today will die prematurely as a result of tobacco use, with 1 billion deaths from tobacco expected during this century.³ More important is the fact that this epidemic of disease and death caused by tobacco is increasing very rapidly.¹

There are almost 275 million tobacco users in India. Among adults over one- third (35%) of population use tobacco products, with 48% males and 20% of females

using some form of tobacco.⁴ In India, tobacco consumption is responsible for half of all the cancers in men and a quarter of all cancers in women,⁵ in addition to being a risk factor for cardiovascular diseases and chronic obstructive pulmonary diseases.⁶ India also has one of the highest rates of oral cancer in the world, partly attributed to high prevalence of tobacco chewing.¹⁰

Many studies⁸⁻¹⁰ have shown that the morbidity and mortality associated with tobacco use is largely preventable by taking measures to prevent the initiation & promote the cessation of tobacco use. However the varied patterns of tobacco use and diversity of socio-economic determinants substantially influence the profile of tobacco related diseases. Data from research studies is required to quantify the problem and identify its determinants which need to be area specific to reduce the burden of tobacco use and provide a credible basis for evolving tobacco control policies.¹ Hence this paper seeks to determine the prevalence of tobacco use in

Address Correspondence to :

Dr. Shital Gaude

Junior Consultant, NCDC, Directorate of Health Services, Campal, Panaji, Goa.

Email : shital19@gmail.com

Mobile : 8806721413



the individuals aged 20 years and above in the urban area of Santa Cruz. It also seeks to determine the prevalence of the various forms of tobacco used in the study population. Some socio-demographic factors associated with tobacco use have also been examined.

Material and methods : The present cross sectional community based study was conducted in the urban area of Santa Cruz in Goa. The study was carried out over a period of 6 months from November 2012 to May 2013 in individuals aged 20 years and above. Sample size for the study was calculated with the formula:

$$N = Z\alpha^2 pq / d^2$$

Where,

N= Sample size, $Z\alpha$ = Confidence level = 1.96, p= Prevalence of Tobacco use in Goa, Q= 1-p, d = Permissible error.

311 households were chosen using systematic random sampling and 935 individuals aged 20 years and above from the 311 households were recruited in the study. The data was collected in a pre-designed structured questionnaire by face to face interview.

Written consent was obtained from all the study subjects. Questionnaire included both open and close ended questions. It consisted of 2 parts:

1. Household Questionnaire and
2. Individual Questionnaire.

The results of the present study were analysed using SPSS version 14 and statistical test (Chi Square test) was used where applicable.

Definitions :

Ever users of tobacco: Ever Users of Tobacco include Current and Former smoking, smokeless and any form tobacco users and was defined as an adult who had smoked 100 cigarettes and/or 20 or more smokeless tobacco forms orally within the six months prior to commencement of the study.¹¹⁻¹⁴

Current Smoking tobacco user : An adult who had smoked 100 smoking forms in his or her lifetime and/or who had smoked within the six months prior to commencement of the study.¹³

Former Smoking tobacco user: Former smoker was defined as an adult who had smoked 100 smoking forms in his/her lifetime and did not smoke during the last six months prior to commencement of the study.¹³

Current Smokeless Tobacco Users : Current smokeless tobacco user was defined as an adult who had used 20 or more smokeless tobacco forms orally within six months prior to commencement of the study.^{12,14}

Former Smokeless Tobacco Users : was defined as an adult who had used 20 or more smokeless tobacco forms

but did not use for the last six months prior to commencement of the study.^{12,14}

Any form (Both forms) : referred to use of both smoking and smokeless forms of tobacco by an adult at any period of time wherein more than 100 smoking forms and more than 20 smokeless forms have been used prior to commencement of the study.¹²⁻¹⁵

Never Users : A never user was defined as one who has never used tobacco in any form, or who had used either less than 100 smoking tobacco forms or less than 20 smokeless tobacco forms in his or her lifetime.^{14,16,17}

Results : In the present study, socio demographic and general characteristic of study population is given in table 1. It reveals highest percentage of males (25.7%) in the age group of 20-29 years and lowest percentage of males (15.6%) in the age group of 60 years and above. Highest percentage of females (25.8%) was found in the age group of 20-29 years and lowest percentage of females (15.4%) was found in the age group of 60 years and above. The mean age of study population was found to be 41.73 ± 15.5 years.

The prevalence of tobacco use was found to be 34.8%. i.e. 326 individuals out of total 935 individuals were found to be using tobacco. The prevalence of smoking tobacco use was found to be 20.3% i.e. 190 individuals out of total 935 were found to be using smoking form of tobacco. The prevalence of smokeless tobacco use was found to be 14.1 i.e. 132 individuals out of total 935 were found to be using smokeless form of tobacco. The prevalence of tobacco use in any form was found to be 0.4% i.e. 4 individuals out of total 935 were found to be using any forms of tobacco.

It was observed that, in males, the highest percentage of ever users of tobacco (40.8%) was found in the age group between 40-49 years of age and lowest (10.7%) in the age group of 20-29 years. In females, the highest percentage of ever users of tobacco (9.1%) was found in the age group between 40-49 years of age and lowest (4.8%) in the age group of 60 years and above. The mean age of ever users of tobacco was 43.31 ± 11.5 years in males and 40.63 ± 12.9 in females.

The prevalence of smoking in males was found to be higher (39.2%) than that in females (0.8%). The prevalence of smokeless tobacco use was also found higher in males (16.6%) than that in females (11.4%).

The commonest smoking tobacco form used was found to be cigarette (92%), followed by Beedi (6%) and Dhumti (2%) and the commonest smokeless tobacco product used was found to be Betel Quid (51%), followed by Gutkha (38%), Toothpaste (6.0%), Mishri (3%) and Khaini (1%).

It was observed that highest percentage of both male and female ever users of tobacco were below 20 years of age at initiation of tobacco. In males mean age at initiation was 18.85 ± 4.3 years and in females the mean age at initiation was 18.19 ± 1.5 years.

It was observed that, the percentage of tobacco use was found to be highest in Muslims (39.5%), followed by Hindus (34.6%) and Catholics (33.6%). The percentage of tobacco use was found to be highest in married people (39.8%), followed by among widow/widower/divorced/separated group (29.0%) and unmarried people (12.7%).

In the present study it was observed that, the percentage of tobacco use was found to be highest in illiterate individuals (40.0%) and lowest in individuals with graduation and above education (21.3%). The percentage of tobacco use was found to be highest in unemployed (65.2%). No student was found to be an ever user of tobacco. The percentage of ever users of tobacco was found to be highest in individuals belonging to lower socio-economic class (51.5%) while lowest in those belonging to upper socio-economic class (17.3%).

In the present study the association between various socio-demographic factors i.e. age, sex, marital status, education, occupation and socio-economic class and tobacco use was found to be statistically significant.

Discussion and Conclusion:

Prevalence of Tobacco Use : In the present study, the prevalence of tobacco use was found to be 34.8%. i.e. 326 individuals out of total 935 individuals were found to be using tobacco. The prevalence of tobacco use in the present study was found to be similar to the prevalence in other studies. Global Adult Tobacco Survey India (GATS India)¹⁹ found the prevalence of 35% in the population aged 15 years and above. In a study conducted by Rani et al in 2003²⁶, it was found that 30 percent of the population aged 15 years and above were tobacco users. Survey conducted by the Indian Council of Medical Research in Karnataka and Uttar Pradesh found the prevalence of tobacco use as 29.6% in Karnataka and 34.6% in Uttar Pradesh¹⁸.

In males, the highest percentage of ever users of tobacco (40.8%) was found in the age group between 40-49 years of age and lowest (10.7%) in the age group of 20-29 years. In females, the highest percentage of ever users of tobacco (9.1%) was found in the age group between 40-49 years of age and lowest (4.8%) in the age group of 60 years and above. The mean age of ever users of tobacco was 43.31 ± 11.5 years in males and 40.63 ± 12.9 in females. Similar findings have been reported in other studies. Rani et al²⁶ reported that the prevalence of tobacco use increased up to the age of 50 years and then

levelled or declined which was similar to the present study. As also Laishram Jenibala Devi et al⁽²⁵⁾ in their study found that the age wise prevalence of tobacco use was higher as the age advanced and the highest rate was found in the age group of 40-49 years and then declined gradually as age advances.

Sex and Tobacco use :

The prevalence of ever users of tobacco was higher in males (56.7%) than that in females (12.3%), which was found to be statistically significant ($\chi^2 = 202.75$; $df = 1$; $p = 0.000$). Similar findings were reported in other studies. In GATS India 2010¹⁹, the prevalence of both smoking and smokeless tobacco use in males was higher than in females. So also, Rani et al²⁶, in their study found that, 47% men and 14% of women either smoked or chewed tobacco. In a survey in rural Nagaur, Rajasthan, 51% of males and 5% of females were tobacco users among 3148 respondents 21 years of age and above²³. In urban Jaipur, in three successive studies about 39% of men and 17% of women 20 years and above were tobacco users²².

Forms of Tobacco used :

In the present study prevalence of smoking tobacco use was found to be 20.3% i.e 190 individuals out of total 935 were found to be using smoking form of tobacco. The prevalence of smokeless tobacco use was found to be 14.1 i.e 132 individuals out of total 935 were found to be using smokeless form of tobacco. The prevalence of tobacco use in any form was found to be 0.4% i.e. 4 individuals out of total 935 were found to be using any forms of tobacco. Similar findings were reported in other studies. In a survey conducted by the Indian Council of Medical Research on prevalence of tobacco use in Karnataka and Uttar Pradesh in India¹⁸, the prevalence of current use of tobacco in smoking, smokeless form and mixed type was observed to be 14.6%, 12.9% and 0.9%, respectively in Karnataka and 16.9%, 16.4% and 1.1% respectively, in Uttar Pradesh. Rani et al²⁶ reported, 16% of the study population (29.3% men and 2.3% women) smoked tobacco; 20% of the study population (28.1% men and 12.0% women) chewed tobacco/pan masala; and 30% of the study population (46.5% men and 13.8% women) either smoked or chewed tobacco.

The highest percentage (28.7%) of current smoking tobacco use was found to be in the age group of 40-49 years and lowest percentage (4.1%) was found to be in the age group of 20-29 years. The highest percentage (13.1%) of former smoking tobacco use was found to be in the individuals 60 years and above. It was also observed that the highest percentage (19.5%) of current smokeless tobacco use was found to be in the age group of 40-49 years and lowest percentage was found to be

(8.2%) in the individuals 60 years and above. Only 1 former smokeless tobacco user in the age group of 30-39 years was reported in the study. The percentage of current tobacco users of any forms was found to be highest (1.3%) in individuals 60 years and above. Similar findings were reported in other studies. Ansari *et al*⁽³⁾ reported that the highest number of smoking tobacco users were in the age group of 40-49 years of age, however highest number of smokeless tobacco users were reported in the age group of 30-39 years of age. H. K. Chaturvedi *et al*⁽²⁴⁾ in their study reported that highest number (35.7%) of current smokers were in the age group of 30-49 years of age and highest number (29.5%) of smokeless tobacco users were also in the age group of 30-49 years of age.

The percentage of current smoking in males was found to be highest (35.9%) in the age group of 40-49 years and the lowest percentage (4.6%) was found in the age group of 20-29 years. The percentage of current smoking in females was found to be highest (1.5%) in the age group of 50-59 years. In males the highest percentage of former smokers was found in the age group of 60 years and above. Only one female was found to be former tobacco user in the age group of 60 years and above. The prevalence of smoking in males was found to be higher (39.2%) than that in females (0.8%). Similar findings were reported in other studies. GATS India⁽¹⁹⁾ found the prevalence of smoking tobacco use was higher (15%) in males than in females (1.9%). Ankur Garg *et al*⁽²⁾ also in their study found the prevalence of smoking in males to be higher (40.1%) than in females (8.8%). The commonest smoking tobacco form used was found to be cigarette (92%), followed by Beedi (6%) and Dhumti (2%). However, in a survey Conducted by the Indian Council of Medical Research on prevalence of Tobacco Use in Karnataka and Uttar Pradesh in India⁽¹⁸⁾, Bidi was observed to be the most popular modality of tobacco smoking accounting for 91.7% and 84.5% of smoking habit (with or without other tobacco habits) in Karnataka and Uttar Pradesh. The difference in the predominant smoking form used in the present study could probably be attributed to the high socio-economic status of the people as well urban culture.

The percentage of current smokeless tobacco use in males was found to be highest (18.0%) in the age group of 40-49 years and the lowest percentage (5.0%) was found in the age group of 20-29 years. The percentage of current smokeless tobacco use in females was found to be highest (3.8%) in the age group of 20-29 years. Only 1 male individual was found to be former smokeless tobacco user in the age group of 60 years and above. No

female former tobacco user was found in the study. The prevalence of smokeless tobacco use in males was found to be higher (16.6%) than that in females (11.4%). Similar findings were reported in other studies. GATS India⁽¹⁹⁾ found that the prevalence of smokeless tobacco use was higher (20.3%) in males than in females (17.3%). Avinash Kumar *et al*⁽⁴⁾ found in their study that the prevalence of smokeless tobacco use in males was higher (76.41%) than that in females (63.70%).

In the study the commonest smokeless tobacco product used was found to be Betel Quid (51%), followed by Gutkha (38%), Toothpaste (6.0%), Mishri (3%) and Khaini (1%). However, Avinash Kumar *et al* in their study found that the khaini was the commonest form of smokeless tobacco used by all the groups. Ghutka was the second most commonly used tobacco among all the groups except business and service class where betel quid was more common. Gupta and Ray, 2003⁽²¹⁾; Dobe *et al.*, 2006⁽¹⁵⁾ reported that smokeless tobacco is consumed predominantly by chewing in form of pan (piper betel leaf filled with sliced areca nut, lime, catechu, and other spices chewed with or without tobacco), pan-masala or gutkha and mishri.

The combinations of Cigarettes and Gutkha, Cigarettes and Betel Quid and Bidi and Betel Quid were used, and only 4 male individuals reported using them. Of the three combinations used cigarette and gutkha was the predominant combination used.

The percentage of tobacco use was found to be highest in Muslims (39.5%), followed by Hindus (34.6%) and Catholics (33.6%). Similarly, Dixit S *et al*⁽²⁷⁾, in their study found high prevalence in Muslims though religion was found to be insignificantly related to tobacco use.

It was observed that, the percentage of tobacco use was found to be highest in married people (39.8%), followed by among widow/widower/divorced/separated group (29.0%) and unmarried people (12.7%). This is attributed to the fact that the sample size of married people is quite higher as compared to both other groups. The variables tobacco use and marital status were found to be associated i.e. there is statistically significant difference in tobacco use among different marital groups. Similarly, T. Rooban *et al*⁽²⁸⁾, in their study on tobacco use among men in urban slums and non slums, found that tobacco use in smokeless, smoking and in any form was highest in married (67.3%, 72.1% and 63.3% respectively) among slum dwellers and (67.9%, 71.1%, and 67.4% respectively) among non slum dwellers. So also S. Dixit *et al*⁽²⁷⁾ reported that tobacco use was highest in married people (Smoking-85.1%, Smokeless- 80%).

It was observed that, the percentage of tobacco use was found to be highest in illiterate individuals (40.0%) and lowest in individuals with graduation and above education (21.3%). The variables tobacco use and education were found to be associated i.e there is statistically significant difference in tobacco use among different educational groups. Similar findings were reported in other studies. Laishram Jenibala Devi *et al*²⁵ in their study found that the educational status was significantly associated with tobacco use. This is consistent with observations that those with lower level of education are more likely to use tobacco. In a survey Conducted by the Indian Council of Medical Research on prevalence of Tobacco Use in Karnataka and Uttar Pradesh in India 2001¹⁸, negative association between education and prevalence of tobacco use was observed overall in both the States. In a study by Basan Gauda Patil¹⁶, the higher prevalence of tobacco use was found among illiterates(35.1%) & lower in postgraduates (15%).

It was observed that, the percentage of tobacco use was found to be highest in unemployed (65.2%). No student was found to be an ever user of tobacco. The variables tobacco use and occupation were found to be associated i.e there is statistically significant difference in tobacco use among different occupational groups. Similarly, GATS India 2010¹⁹ found that amongst all occupational categories tobacco use is lowest among students. B. Prabhakar *et al*⁵ reported that the prevalence of current tobacco use was seen to be the lowest among students for both men and women. Also in a prospective cohort study conducted in Mumbai²⁰, unskilled male workers, male service workers and unemployed individuals were more at risk than professionals.

It was observed that, the percentage of ever users of tobacco was found to be highest in individuals belonging to lower socio-economic class (51.5%) while lowest in those belonging to upper socio-economic class (17.3%). The variables tobacco use and socioeconomic status were found to be associated i.e. there is statistically significant difference in tobacco use among different socioeconomic classes. Similar findings were reported in other studies. In a study by Basan Gauda Patil¹⁶, the prevalence in class I was 23.6% and it increased steadily across group to reach a maximum level of 37.5% in class V. In a survey conducted by the Indian Council of Medical Research on prevalence of Tobacco Use in Karnataka and Uttar Pradesh in India 2001¹⁸, it was found that higher family income levels were associated with a lesser prevalence of current tobacco use.

Conclusion :

The high prevalence of tobacco use among men and women points towards the fact that mere knowledge about the health hazards is not sufficient to make them stop using tobacco. So there is a need to develop effective health education and multi-factorial tobacco quitting strategies with focus on help and support for those who wish to quit tobacco.




Fig 1: Pie chart showing distribution of smoking forms of tobacco




Fig 2: Pie chart showing distribution of smokeless forms of tobacco




Fig 3: Pie chart showing distribution of any forms of tobacco

TABLE 1: Socio demographic profile and general characteristic of study population.

Socio-demographic and general profile	Male		Female		Total	
	No	%	No	%	No	%
Age						
20-29	122	25.7	119	25.8	241	25.8
30-39	113	23.8	111	24.0	224	24.0
40-49	88	18.5	86	18.6	174	18.6
50-59	77	16.2	74	16.0	151	16.3
>=60	74	15.6	71	15.4	145	15.5
Religion						
Hindu	360	59.1	249	40.9	609	65.1
Catholic	168	71.5	67	28.5	235	25.1
Muslim	55	60.4	36	39.6	91	9.8
Marital Status						
Unmarried	78	58.2	56	41.8	134	14.3
Married	365	51.6	343	48.4	708	75.7
Widow/Widower/Divorced/Separated	19	20.4	74	79.6	93	9.9
Levels of Education						
Illiterate	71	33.0	144	67.0	215	22.9
Primary	66	47.1	74	52.9	140	14.9
Secondary	144	53.7	124	46.3	268	28.6
Higher Secondary	77	55.8	61	44.2	138	14.7
Graduation and Above	104	59.8	70	40.2	174	18.6
Occupation						
Unemployed	52	36.9	89	63.1	141	15.0
Unskilled	92	53.8	79	46.2	171	18.2
Skilled	114	78.1	32	21.9	146	15.6
Business	234	91.8	21	8.2	255	27.2
Professional	121	71.2	49	28.8	170	18.1
Student	25	48.1	27	51.9	52	5.5
Socio-Economic class						
Upper class	50	51.0	48	49.0	98	10.4
Upper middle	125	51.0	120	49.0	245	26.2
Lower middle	116	51.3	110	48.6	226	24.1
Upper lower	86	49.4	88	50.6	174	18.6
Lower	99	51.6	93	48.4	192	20.5

TABLE 2: Distribution of tobacco use according to age and sex.

Age Group	Ever Users (Tobacco)		Never Users		Total
	Male	Female	Male	Female	
20-29	26(10.7%)	12(4.9%)	96(39.8%)	107(44.3%)	241(100.0%)
30-39	61(27.2%)	13(5.8%)	52(23.2%)	98(43.7%)	224(100.0%)
40-49	71(40.8%)	16(9.1%)	17(9.7%)	70(40.2%)	174(100.0%)
50-59	60(39.7%)	9(5.9%)	17(11.2%)	65(43.0%)	151(100.0%)
>=60	51(35.1%)	7(4.8%)	23(15.8%)	64(44.1%)	145(100.0%)
Total	269(28.7%)	57(6.0%)	205(21.9%)	404(43.2%)	935(100%)

(Figures in parenthesis indicate percentage)

TABLE 3: Distribution of forms of tobacco used according to age.

Age Group	Tobacco Forms							Total
	Smoking		Smokeless		Any forms		Never	
	Current	Former	Current	Former	Current	Former	User	
20-29	10 (4.1%)	0	28 (11.6%)	0	0	0	203 (84.2%)	241 (100.0%)
30-39	39 (17.4%)	0	34 (15.1%)	1 (0.4%)	0	0	150 (66.9%)	224 (100.0%)
40-49	50 (28.7%)	1 (0.5%)	34 (19.5%)	0	1 (0.5%)	0	87 (50.0%)	174 (100.0%)
50-59	42 (27.8%)	4 (2.6%)	23 (15.2%)	0	1 (0.6%)	0	82 (54.3%)	151 (100.0%)
>=60	25 (17.2%)	19 (13.1%)	12 (8.2%)	0	2 (1.3%)	0	87 (60%)	145 (100.0%)
Total	165 (17.6%)	24 (2.5%)	78 (8.3%)	1 (0.1%)	4 (0.4%)	0	609 (65.1%)	935 (100%)

(Figures in parenthesis indicate percentage)

TABLE 4: Socio-demographic factors and Tobacco use

	Ever users (Tobacco)		P value
	Number	%	
Religion			
Hindu	211	34.6	p>0.05
Catholic	79	33.6	
Muslims	36	39.5	
Marital Status			
Unmarried	17	12.7	$\chi^2 = 38.10$; df=2; p=0.000
Married	282	39.8	
Widow/Widower/Divorced/Separated	27	29.0	
Levels of Education			
Illiterate	86	40.0	$\chi^2 = 20.56$; df=4; p=0.001
Primary	51	36.4	
Secondary	105	39.1	
Higher Secondary	47	34.1	
Graduation and Above	37	21.3	
Occupation			
Unemployed	92	65.2	$\chi^2 = 162.8$; df=7; p=0.000
Unskilled	80	47.0	
Skilled	41	28.1	
Business	76	44.4	
Professional	37	14.5	
Student	0	-	
Socio-Economic class			
Upper class	17	17.3	$\chi^2 = 16.28$; df=4; p=0.003
Upper middle	81	33.0	
Lower middle	76	33.6	
Upper lower	53	30.4	
Lower	99	51.5	

References:

1. Anantha N, Nandakumar A, Vishwanath N, Venkatesh T, Pallad YG, Manjunath P, et al. Efficacy of an antitobacco community program in India. *Cancer Causes and Control* 1995; 6:119-29.
2. Ankur Garg. Prevalence and correlates of tobacco smoking, awareness of hazards, and quitting behavior among persons aged 30 years or above in a resettlement colony of Delhi, India, 2012;29:336-340.
3. Ansari MM, Beg MH, Haleem S. Clinicopathological profile of carcinoma of oesophagus at Aligarh. *J Indian Med Assoc.* 1991;89(8):217-9.
4. Avinash et al. A study on knowledge, attitude and practices regarding smokeless tobacco use among adult (more than eighteen years) in the rural area of Jharkhand. *Int J Basic App Med Sci.* 2013; 3:378-81.
5. SS Narake, MS Pednekar. Social disparities in tobacco use in India: The roles of occupation, education and gender. *Indian Journal of Cancer:* 2012; 49: 401-9.
6. Barbeau EM, Krieger N, Soobader M. Working class matters: socioeconomic disadvantage, race/ethnicity, gender, and smoking in NHIS 2000. *Am J Public Health* 2004; 94:269–78.
7. Bhonsle RB, Murti PR, Gupta PC. Tobacco habits in India. In: Gupta PC, Hamner JE, Murti PR. *Control of tobacco-related cancers and other diseases.* International Symposium 1990. Bombay: Oxford University Press; 1992:25-45.
8. Center for Disease Control and Prevention, Health Effects of Cigarette Smoking. http://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/Health_Effects_of_Cigarette_Smoking. Obtained July 22, 2009.
9. Chavan BS, Arun P et al. Prevalence of alcohol & drug dependence in rural & slum population of Chandigarh: A community survey. *Indian Journal of Psychiatry.* 2007;49(1):44-8.
10. Chandra V, Ganguli M. Smoking among the elderly in rural Haryana (India): Khaini. New Delhi: WHO SEARO; 2002.
11. De Silva V, Samarasinghe D, Hanwella R. Et al. Association between concurrent alcohol and tobacco use and poverty. *Drug Alcohol Rev.* 2011;30(1):69-73.
12. Difranza, J.R., and Guerrera, M.P. Alcoholism and smoking. *Journal of Studies on Alcohol* 1990; 51(2): 130–5.
13. Dobe M, Sinha DN, Rahman K. Smokeless tobacco use and its implications in WHO South East Asia Region. *Indian J Public Health* 2006;50:70-5.
14. Donna L. Ansara et al. Tobacco Use by Men and Women in 49 Countries with Demographic and Health Surveys. 2013.
15. Douglas E. Jorenby. The Merck Manual- Home Health Handbook. Last modified in May 2013.
16. Dr. Basanagauda K. Patil. A community based study on tobacco use in Davangere Taluka, Karnataka.
17. Dr. K. Venugopal. New methods and drugs to stop smoking. Why, stubb out lives? Kerala Calling July 2004.
18. Dr. Kishore Chaudhry Deputy Director General Indian Council of Medical Research Prevalence of Tobacco Use in Karnataka and Uttar Pradesh in India 2001.2
19. Global Adult Tobacco Survey India Report, 2009-2010.
20. Glorian Sorensen et al. Social Disparities in Tobacco Use in Mumbai, India: The Roles of Occupation, Education, and Gender. *Amer J Pub Health.* 2005; 96 (6):1003-8.
21. Gupta PC, Ray CS. Smokeless tobacco and health in India and South Asia. *Respirology* 2003;8:419-31.
22. Gupta R, Gupta VP, Sarna M, Bhatnagar S, Thanvi J, Sharma V, et al. Prevalence of coronary heart disease and risk factors in an urban Indian population: Jaipur Heart Watch-2. *Ind Heart Journal* 2002; 54: 59;66.
23. Gupta R, Prakash H, Majumdar S, Sharma S, Gupta VP. Prevalence of coronary heart disease and coronary risk factors in an urban population of Rajasthan. *Ind Heart Journal* 1995; 47:331.8.
24. H.K. Chaturvedi. Tobacco use in Mizoram, India: Sociodemographic differences in patterns. *Southeast Asian J Trop Med Pub Health.* 1998; 29 (1): 66-70.
25. Laishram Jenibala Devi, Jalina Laishram, Laishram Jayarani, Dilip Ingudum, H Sanayaima Devi. "Prevalence and determinants of tobacco use among adults in an urban community of Imphal west, Manipur". *J Evol Med Dental Sci* 2013; 12;5302-8.
26. M Rani, S Bonu, P Jha, S N Nguyen, L Jamjoum. Tobacco use in India: prevalence and predictors of smoking and chewing in a national cross-sectional household survey. *tobaccocontrol.bmj.com* 2003 12: e4.
27. S. Dixit, M. A. Ansari, Z. Khan, N. Khaliq. Prevalence and predictors of tobacco use; a cross-sectional household survey in aligarh district of Uttar Pradesh. *Indian Journal of Community Health* 2012; 24 (3): 203-08.
28. T. Rooban et al. Prevalence of tobacco use among urban adult men in India: A comparison of slum dwellers v/s non-slum dwellers. *Indian J Dental Research* 2012; 23 (1): 31-8.

How to Cite this article :

Rama G S, Vanita S P, Prevalence and patterns of smoking and smokeless tobacco use in Santa-Cruz, Goa, India. *J Pub Health Med Res*, 2015;3(1):21-28

Funding: Declared none

Conflict of interest: Declared none